

2215/14

SE/EXTC / III CBGS  
A.M. III

QP Code : NP-18646

(3 Hours)

[ Total Marks : 80

- N. B. : (1) Question No. 1 (one) is compulsory.  
 (2) Attempt any 3 (three) questions from the remaining questions.  
 (3) Assume suitable data, if necessary.

1. (a) Evaluate  $\int_0^{\infty} \frac{(\cos 6t - \cos 4t)}{t} dt$  5  
 (b) Obtain complex form of fourier series for  $f(x) = e^{ax}$  in  $(-1,1)$  5  
 (c) Find the work done in moving a particle in a force field given by  $\vec{F} = 3xy\hat{i} - 5z\hat{j} + 10x\hat{k}$  along the curve  $x = t^2 + 1$ ,  $y = 2t^2$ ,  $z = t^3$  from  $t = 1$  to  $t = 2$ . 5  
 (d) Find the orthogonal trajectory of the curves  $3x^2y + 2x^2 - y^3 - 2y^2 = \alpha$ , where  $\alpha$  is a constant. 5
2. (a) Evaluate  $\frac{d^2y}{dt^2} + 2\frac{dy}{dt} - 3y = \sin t$ ,  $y(0) = 0$ ,  $y'(0) = 0$ , by Laplace transform 6  
 (b) Show that  $J_{5/2} = \sqrt{\frac{2}{\pi x}} \left[ \frac{3-x^2}{x^2} \sin x - \frac{3}{x} \cos x \right]$  6  
 (c) (i) Find the constants a, b, c so that  $\vec{F} = (x + 2y + az)\hat{i} + (bx - 3y - z)\hat{j} + (4x + (y + 2z))\hat{k}$  is irrotational. 4  
 (ii) Prove that the angle between two surfaces  $x^2 + y^2 + z^2 = 9$  and  $x^2 + y^2 - z = 3$  at the point  $(2, -1, 2)$  is  $\cos^{-1}\left(\frac{8}{3\sqrt{21}}\right)$  4
3. (a) Obtain the fourier series of  $f(x)$  given by  $f(x) = \begin{cases} 0 & , -\pi \leq x \leq 0 \\ x^2 & 0 \leq x \leq \pi \end{cases}$  6  
 (b) Find the analytic function  $f(z) = u + iv$  where  $u = r^2 \cos 2\theta - r \cos \theta + 2$  6  
 (c) Find Laplace transform of  
 (i)  $te^{-3t} \cos 2t \cdot \cos 3t$  8  
 (ii)  $\frac{d}{dt} \left[ \frac{\sin 3t}{t} \right]$

Con. 11456-14.

[ TURN OVER

4. (a) Evaluate  $\int J_3(x) dx$  and Express the result in terms of  $J_0$  and  $J_1$  6  
 (b) Find half range sine series for 6  
 $f(x) = \pi x - x^2$  in  $(0, \pi)$

Hence deduce that  $\frac{\pi^3}{32} = \frac{1}{12} - \frac{1}{3^2} + \frac{1}{5^2} - \frac{1}{7^2} + \dots$

- (c) Find inverse Laplace transform of 8

(i)  $\frac{1}{s} \tanh^{-1}(s)$  (ii)  $\frac{se^{-2s}}{(s^2 + 2s + 2)}$

5. (a) Under the transformation  $w + 2i = z + \frac{1}{z}$ , show that the map of the circle  $|z| = 2$  is an ellipse in  $w$ -plane. 6

- (b) Find half range cosine series of  $f(x) = \sin x$  in  $0 \leq x \leq \pi$ . 6  
 Hence deduce that

$$\frac{1}{1.3} + \frac{1}{3.5} + \frac{1}{5.7} + \dots = \frac{1}{2}$$

- (c) Verify Green's theorem, for 8

$\oint_c (3x^2 - 8y^2) dx + (4y - 6xy) dy$  where  $c$  is boundary of the region defined by  $x=0$ ,  $y=0$ , and  $x+y = 1$ .

6. (a) Using convolution theorem; evaluate 6

$$L^{-1} \left\{ \frac{1}{(s-1)(s^2+4)} \right\}$$

- (b) Find the bilinear transformation which maps the points 6  
 $z = 1, i, -1$  onto  $w = 0, 1, \infty$

- (c) By using the appropriate theorem, Evaluate the following :- 8

(i)  $\int \bar{F} \cdot d\bar{r}$  where  $\bar{F} = (2x - y)\hat{i} - (yz^2)\hat{j} - (y^2z)\hat{k}$

and  $c$  is the boundary of the upper half of the sphere  $x^2 + y^2 + z^2 = 4$

(ii)  $\iiint_s (9x\hat{i} + 6y\hat{j} - 10z\hat{k}) \cdot d\bar{s}$  where  $s$  is

the surface of sphere with radius 2 units.